

Mathematics 30–2 Formula Sheet

Relations and Functions

Graphing Calculator Window Format

$$x: [x_{\min}, x_{\max}, x_{\text{scl}}]$$

$$y: [y_{\min}, y_{\max}, y_{\text{scl}}]$$

Exponents and Logarithms

$$y = a^x \leftrightarrow x = \log_a y$$

$$\log_b c = \frac{\log_a c}{\log_a b}$$

Laws of Logarithms

$$\log_a(M \cdot N) = \log_a M + \log_a N$$

$$\log_a\left(\frac{M}{N}\right) = \log_a M - \log_a N$$

$$\log_a(M^n) = n \log_a M$$

Exponential functions

$$y = a \cdot b^x$$

Sinusoidal functions

$$y = a \cdot \sin(bx + c) + d$$

$$\text{Period} = \frac{2\pi}{b}$$

Quadratic equations

$$\text{For } ax^2 + bx + c = 0$$
$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Probability

$$n! = n(n-1)(n-2)\dots 3 \cdot 2 \cdot 1$$

where $n \in N$ and $0! = 1$

$${}_n P_r = \frac{n!}{(n-r)!}$$

$${}_n C_r = \frac{n!}{(n-r)!r!}$$

$${}_n C_r = \binom{n}{r}$$

$$P(A \cup B) = P(A) + P(B)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A \cap B) = P(A) \cdot P(B|A)$$

Logical Reasoning

A' Complement

\emptyset Empty set

\cap Intersection

\subset Subset

\cup Union